

# Calcium Dependence of Settlement and Nematocyst Discharge in Actinulae of the Hydroid *Tubularia mesembryanthemum*

SATORU KAWAI<sup>1</sup>, KEIJI YAMASHITA<sup>2</sup>, MITSUYO NAKAI, MIYUKI TAKAHASHI,  
AND NOBUHIRO FUSETANI<sup>3,\*</sup>

*Biofouling Project, ERATO, Research Development Corporation of Japan, c/o Yokohama R&D center,  
Niigata Engineering Co. Ltd., 27 Shin-isogo-cho, Isogo-ku, Yokohama, 235, Japan*

**Abstract.** The influence of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions on both *atrachous isorhiza* (AI) discharge and settlement of actinular larvae of the hydroid *Tubularia mesembryanthemum* was investigated.  $\text{Mg}^{2+}$ -supplemented artificial seawater (ASW) completely inhibited both events at a concentration of 206 mM, whereas lowered  $\text{Mg}^{2+}$  concentrations enhanced them.  $\text{Ca}^{2+}$  ions in the bathing solution highly regulated AI discharge and settlement, and  $\text{Mg}^{2+}$  ions may down-regulate these events. The effect of inorganic  $\text{Ca}^{2+}$ -channel blockers, including  $\text{Gd}^{3+}$  and  $\text{La}^{3+}$ , was also examined. Larval settlement was inhibited by  $\text{Co}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{La}^{3+}$ , and  $\text{Gd}^{3+}$ , with half inhibitory concentrations ( $\text{IC}_{50}$ ) of 5800, 260, 53, 45, and 7  $\mu\text{M}$ , respectively; AI discharge was also inhibited by these ions, with  $\text{IC}_{50}$  values of 6600, 500, 78, 41, and 5  $\mu\text{M}$ , respectively. These results suggest possible involvement of stretch-activated  $\text{Ca}^{2+}$  channels in the signal transmission of both AI discharge and larval settlement.

## Introduction

The colonial marine hydroid *Tubularia mesembryanthemum* Allman, 1871 (Hydrozoa: Tubulariidae) is a prolific marine fouling organism that frequently dominates artificial substrata used in aquaculture. Actinula larvae, produced as

a means of dispersal, are composed of a conical manubrium, oral- and aboral tentacles, and a basal protrusion that later becomes the settlement organ. Settlement-competent actinulae initiate settlement behavior by contact of the aboral tentacle tip (ATT) with the substratum surface, which triggers discharge of *atrachous isorhiza* (AI), the agglutinant nematocysts located at the ATT.

Upon reception of appropriate stimuli, nematocysts can eject adhesive tubules (Tardent, 1988). Nematocyte excitation requires both mechanical and chemical stimuli (Pantin, 1942). Two classes of chemoreceptors were identified in the tentacle of the sea anemone *Aiptasia pallida*: one sensitive to low molecular weight amino/imino compounds, and the other specific for N-acetylated sugars (Thorington and Hessinger, 1988a). In addition, two types of mechanoreceptors are involved in triggering nematocyst discharge; one is contact-sensitive (Thorington and Hessinger, 1988b) and the other is tunable, vibration- and frequency-sensitive (Watson and Hessinger, 1989, 1991).

The necessity of external  $\text{Ca}^{2+}$  ions for nematocyst discharge has been demonstrated in *Aiptasia mutabilis* (Santoro and Salleo, 1991b), *Pelagia noctiluca* (Salleo *et al.*, 1994a), *Calliactis parasitica* (Salleo *et al.*, 1994b), *Charybdea rastonii* (Yanagita, 1973), and *Anthopleura elegantissima* (McKay and Anderson, 1988). In *Hydra vulgaris*, discharge of *stenotele* is regulated by a voltage- and  $\text{Ca}^{2+}$ -dependent mechanism, allowing  $\text{Ca}^{2+}$  influx from the bathing solution (Gitter *et al.*, 1994). The involvement of intracellular  $\text{Ca}^{2+}$  ions in the metamorphosis of hydrozoan larvae has also been suggested on the basis of studies monitoring the emission of bioluminescence from photoproteins, thereby visualizing changes in  $[\text{Ca}^{2+}]_i$  during

Received 3 February 1998; accepted 19 October 1998.

<sup>1</sup> Present address: Natl. Inst. Fruit Sci., Min. Agric. Forest. Fish., Okitsu, Shimizu, Shizuoka 424-0204, Japan. E-mail: kw4682@okl.affrc.go.jp

<sup>2</sup> Present address: Himeji Kisho Co. Ltd., 1417-1 Kiba, Himeji-shi, Hyogo 672, Japan.

<sup>3</sup> Present address: Laboratory of Aquatic Natural Products Chemistry, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113, Japan.

\* Author to whom correspondence should be addressed.

Abbreviations: AI, *atrachous isorhiza*; ATT, aboral tentacle tip.

planular metamorphosis (Freeman and Ridgway, 1987, 1990).

Previous work has shown that  $\text{Ca}^{2+}$  release from intracellular stores triggered AI discharge in *Tubularia mesembryanthemum* actinulae, followed by the inflow of  $\text{Ca}^{2+}$  ions from the bathing solution into the nematocysts (Kawaii *et al.*, 1997). Furthermore, AI discharge was usually accompanied by sinuous movement of the aboral tentacle. The involvement of  $\text{Ca}^{2+}$ -mediated signal transduction in actinular settlement and metamorphosis was suspected. To clarify the role of  $\text{Ca}^{2+}$  ions in actinular settlement, we examined AI discharge and larval settlement using intact larvae that were bathed in artificial seawater (ASW) containing various concentrations of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions. The influence of di- and trivalent cations and  $\text{Ca}^{2+}$ -channel blockers (Yang and Sachs, 1989; Santoro and Salleo, 1991; Salleo *et al.*, 1994a, b; Gitter *et al.*, 1994) was also examined.

## Materials and Methods

### Reagents

Reagent grade artificial seawater (ASW) consisting of NaCl (460 mM), KCl (10.1 mM),  $\text{CaCl}_2$  (9.2 mM), and  $\text{MgCl}_2$  (42.6 mM) at pH 7.6 was adjusted by addition of 5 mM imidazole. The concentration of  $\text{Ca}^{2+}$  was reduced, without altering the osmolarity of the ASW, by adjusting  $\text{Mg}^{2+}$  accordingly.  $\text{Mg}^{2+}$ -supplemented ASW with 59, 75, 125, and 206 mM of  $\text{Mg}^{2+}$  was prepared by mixing the regular ASW with 370 mM  $\text{MgCl}_2$  aqueous solution (which is isotonic to ASW) in the ratios of 1:19, 1:9, 1:3, and 1:1, respectively, to compensate for the osmolarity. In the case of  $\text{Ca}^{2+}$ -channel blockers, the osmolarity was adjusted by  $\text{Na}^+$ . All  $\text{Ca}^{2+}$ -channel blockers were purchased from Wako Pure Chemical Industries (Osaka, Japan).

### Biological materials

Mature *Tubularia mesembryanthemum* colonies were collected mainly from submerged fisheries nets and ropes in the vicinity of Nagai Port in Sagami Bay (eastern Japan, 139°30'E, 31°70'N). Colonies were divided into male and female, and were maintained separately with sand-filtered running seawater at  $16 \pm 2^\circ\text{C}$  as described by Yamashita *et al.* (1997). After fertilization, branches of female colonies with polyps bearing many actinulae were placed in filtered seawater (FSW; pore size, 0.45  $\mu\text{m}$ ), and sinking actinulae were collected. A single polyp could produce up to 300 actinulae, which were maintained at  $4^\circ\text{C}$  prior to use in an experiment. The actinulae were transferred to an ASW-containing beaker and maintained at  $21 \pm 2^\circ\text{C}$  for several hours to recover normal responsiveness. Larval age was defined as time following release from the maternal gono-

phore, and 24-h-old actinulae were used in the following experiments unless otherwise stated.

### Settlement assay

Actinular settlement was assayed using six-well polystyrene plates (Corning Cell Wells, Corning, NY), each well containing 6 ml of ASW and 5 actinulae. The plates were placed on an orbital shaker at  $21 \pm 2^\circ\text{C}$ . The number of settled actinulae was counted under a binocular dissection microscope after 30 h. Each experiment was performed in triplicate.

To determine the reversibility of actinular settlement upon replacement of the test ASW with regular ASW, larvae that were largely sinking or attaching to the bottom of the well were washed by several replacements of the surface of the test ASW with regular ASW (roughly half the volume of the test ASW was exchanged with each replacement).

### Larval behavior and atrichous isorhiza discharge

Each actinula was independently examined for the effect of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , or  $\text{Ca}^{2+}$ -channel blockers on its behavior when an ATT was contacted with a clean micropipette or on AI discharge. Actinulae in regular ASW were washed and suspended in test ASWs containing various concentrations of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , or  $\text{Ca}^{2+}$ -channel blockers. Each actinula was held by suction applied through a micropipette attached to the side of the adhesive protrusion. Behavioral responses to ATT contact with a clean micropipette were examined on suction-held actinula whose ATTs did not contact any substrata.

For monitoring AI discharge, an ATT of a suction-held actinula was immobilized and attached to the bottom of a petri dish by applying gentle suction through a second micropipette attached to the "wrist," which is the part between an ATT and an aboral tentacle. AI discharge of the immobilized ATT was then triggered by the addition of 100  $\mu\text{l}$  of ASW that contained 200 mM  $\text{K}^+$  ions ( $\text{K}^+$ -ASW) and had been adjusted to the osmolarity of regular ASW by reducing the  $\text{Na}^+$  concentration. AI discharge was observed through the  $40\times$  objective of a microscope, and the number of AI discharged before the  $\text{K}^+$ -ASW application (usually 0–1) was subtracted from the final total.

To avoid larval disintegration in  $\text{Ca}^{2+}$ -free ASW, a micropipette-held actinula in regular ASW was quickly and thoroughly washed and suspended in  $\text{Ca}^{2+}$ -free ASW. Immediately afterward, an ATT of the actinula was immobilized as described above and AI discharge was triggered.

### $[\text{Ca}^{2+}]_i$ measurement

$[\text{Ca}^{2+}]_i$  was measured in whole mounts of living actinulae ATT in which AI discharge was induced. The actinulae had been treated with the  $\text{Ca}^{2+}$ -chelating fluorescent indi-

cator fura-2 (Kawaii *et al.*, 1997). The fluorescence intensities of ATTs from fura-2-loaded actinulae were approximately 100 times stronger than those from non-labeled actinulae, and were strong enough for [Ca<sup>2+</sup>]<sub>i</sub> measurement. The value of the fluorescence ratio between excitation at 340 nm and 380 nm (*R*<sub>340/380</sub>) was used to indicate [Ca<sup>2+</sup>]<sub>i</sub>. The minimum interval was 1.9 s for ratiometric imaging.

**Results**

*Ca<sup>2+</sup> dependence of actinular settlement*

Reducing the external Ca<sup>2+</sup> concentration had a significant effect on actinular settlement, and the effect was competitively antagonized by Mg<sup>2+</sup> (Fig. 1A). Settlement was comparable to normal in 4.6–9.2 mM Ca<sup>2+</sup>, but it was inhibited at lower concentrations (70% ± 31% at 2.3 mM Ca<sup>2+</sup>, 8% ± 10% at 1.1 mM Ca<sup>2+</sup>).


Raising the Mg<sup>2+</sup> concentration inhibited actinular settlement: 100% settlement was obtained at 75 mM Mg<sup>2+</sup>, 50% at 125 mM Mg<sup>2+</sup>, and 6.7% ± 11.6% at 206 mM.

The relative inhibitory effect of Mg<sup>2+</sup> on actinular settlement was magnified at lower Ca<sup>2+</sup>. In ASW containing 2.3 mM Ca<sup>2+</sup>, settlement was reduced to 53% ± 12% at 59 mM Mg<sup>2+</sup> and completely inhibited at 206 mM Mg<sup>2+</sup>. The effects of Ca<sup>2+</sup> and Mg<sup>2+</sup> concentration on settlement were statistically significant as assessed by two-way ANOVA (Table 1). Furthermore, there was a significant interaction between Ca<sup>2+</sup> and Mg<sup>2+</sup>. In each case the dose-dependent reduction of actinular settlement was reversed by rinsing the larvae with regular ASW 9 h after the experiment.

*Antagonistic effect of Mg<sup>2+</sup>*

Raising the Mg<sup>2+</sup> concentration in an otherwise regular ASW had an inhibitory effect on AI discharge (Fig. 1B); the number of discharged AI was 4.0 ± 4.2 at 125 mM Mg<sup>2+</sup>, and AI discharge was completely arrested at 206 mM Mg<sup>2+</sup>. The inhibitory effect of Mg<sup>2+</sup> increased when the external Ca<sup>2+</sup> ion concentration was lowered. In ASW containing 4.6 mM Ca<sup>2+</sup>, AI discharge was reduced to 5.7 ± 3.8 at 59 mM Mg<sup>2+</sup> and 4.4 ± 3.1 at 75 mM Mg<sup>2+</sup>. In ASW containing 2.3 mM Ca<sup>2+</sup>, discharge was lowered to 4.3 ± 1.4 at 59 mM Mg<sup>2+</sup>, and was completely inhibited at 125 mM Mg<sup>2+</sup>. Two-way ANOVA revealed that the effects of Ca<sup>2+</sup> and Mg<sup>2+</sup> concentration on AI discharge were statistically significant, and that there was a significant interaction between Ca<sup>2+</sup> and Mg<sup>2+</sup> (Table 1). Again, this dose-dependent reduction of AI discharge was reversed by rinsing the larvae with regular ASW 3 h after the experiment.

Figure 2 indicates significant enhancement of AI discharge in Mg<sup>2+</sup>-reduced ASW (one-way ANOVA; *F* = 11.9759, *P* < 0.0001). In this experiment, AI discharge was triggered by K<sup>+</sup>-ASW (100 mM). The number of discharged AI was 2.1 ± 0.6 at 50 mM Mg<sup>2+</sup> (regular ASW)



**Figure 1.** (A) Actinula settlement in artificial seawater with different concentrations of calcium and magnesium. Five larvae were incubated at specified concentrations of external Ca<sup>2+</sup>. Vertical bars indicate standard deviations (*n* = 5). (B) K<sup>+</sup>-ASW triggered *atricous isorhiza* (AI) discharge in artificial seawater containing different concentrations of Ca<sup>2+</sup> and Mg<sup>2+</sup> ions. Means were obtained by averaging the number of discharged AI in each aboral tentacle tip (*n* = 10). Vertical bars indicate standard deviations.

and increased to 7.6 ± 1.6 and 9.2 ± 1.9 at 25 and 10 mM, respectively. Addition of Mg<sup>2+</sup>-reduced ASW (5 mM) also induced sinuous movement of the aboral tentacles.

*Effects of Ca<sup>2+</sup>-channel blockers on actinular settlement*

The effects of inorganic Ca<sup>2+</sup>-channel blockers on actinular settlement were examined in various concentrations of chloride salt cations added to regular ASW. In each case, a dose-dependent reduction of actinular settlement was observed (Fig. 3A). The following ions are listed in order of increasing inhibition efficiency: Co<sup>2+</sup> < Ni<sup>2+</sup> < Cd<sup>2+</sup> < La<sup>3+</sup> < Gd<sup>3+</sup>.

Table 1

Results of two-way ANOVA to assess the effects of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations on (A) actinular settlement and (B) atrichous isorhiza (AI) discharge

Source*	df	F-ratio	P
(A) settlement			
$\text{Ca}^{3+}$	3	118.4902	<0.0001
$\text{Mg}^{2+}$	4	75.7353	<0.0001
$\text{Ca}^{2+}\text{-Mg}^{2+}$	12	9.2255	<0.0001
(B) AI discharge			
$\text{Ca}^{2+}$	3	1022.1822	<0.0001
$\text{Mg}^{3+}$	4	1012.5841	<0.0001
$\text{Ca}^{2+}\text{-Mg}^{2+}$	12	534.8241	<0.0001

\*  $\text{Ca}^{2+}\text{-Mg}^{2+}$  represents the interaction between  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations.

The 50% inhibition concentrations ( $\text{IC}_{50}$ ) were approximately 5800, 260, 53, 45, and 7  $\mu\text{M}$ , respectively. Observed settlement percentages were similar to controls when the blocker was removed 5 h after the experiment.

$\text{K}^+$ -induced AI discharge was also inhibited by these inorganic  $\text{Ca}^{2+}$ -channel blockers (Fig. 3B).  $\text{IC}_{50}$  values for  $\text{Gd}^{3+}$ ,  $\text{La}^{3+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Ni}^{2+}$ , and  $\text{Co}^{2+}$ , were approximately 5, 41, 78, 500, and 6600  $\mu\text{M}$ , respectively. When  $10^4 \mu\text{M}$   $\text{Co}^{2+}$ ,  $10^3 \mu\text{M}$   $\text{Ni}^{2+}$ , and  $10^2 \mu\text{M}$   $\text{Cd}^{2+}$  were added to ASW, AI discharges were lowered to  $1.0 \pm 0.9$ ,  $0.8 \pm 1.2$ , and  $0.6 \pm 1.1$ , respectively. AI discharges were similar to controls when the blocker




Figure 2. Enhancement of atrichous isorhiza (AI) discharge and settlement by low  $\text{Mg}^{2+}$  artificial seawater (ASW). Actinulae were bathed in ASW containing various concentrations of  $\text{Mg}^{2+}$  (regular ASW = 42.6 mM  $\text{Mg}^{2+}$ ) for 1 h, and AI discharge was triggered by  $\text{K}^+$ -ASW (50, 100, or 200 mM) treatment. Means were obtained by averaging the number of discharged AI in each ATT ( $n = 10$ ). Vertical bars indicate standard deviations. Asterisks (\*) indicate significant differences relative to the untreated controls at  $P < 0.05$  by the Tukey-Kramer honestly significant difference test.






Figure 3. Effects of  $\text{Ca}^{2+}$ -channel blockers on actinular settlement (A) and atrichous isorhiza (AI) discharge (B). All experiments were performed using sulfate-free artificial seawater (ASW) at normal pH. Five larvae were incubated for 30 h in ASW containing various concentrations of  $\text{Ca}^{2+}$ -channel blockers.

was removed prior to  $\text{K}^+$ -ASW stimulation 1 h after onset of the experiment. ASW containing 25  $\mu\text{M}$   $\text{Gd}^{3+}$  or 100  $\mu\text{M}$   $\text{La}^{3+}$  decreased the AI discharge to  $1.2 \pm 0.8$  and  $0.8 \pm 1.1$ , respectively, but the discharge was not restored completely when the blockers were removed.

Squashing an ATT with a micropipette caused AI discharge even in the presence of 20  $\mu\text{M}$   $\text{Gd}^{3+}$ , but sinuous tentacle movement, which was usually observed in regular ASW, was inhibited.

#### $[\text{Ca}^{2+}]_i$ transients during atrichous isorhiza discharge

Figure 4 compares representative changes of  $R_{340/380}$  of ATT during AI discharge in  $\text{Ca}^{2+}$ -free ASW,  $\text{Mg}^{2+}$ -supple-



**Figure 4.** [Ca<sup>2+</sup>]<sub>i</sub> transients during *atricous isorhiza* discharge in (A) Ca<sup>2+</sup>-free artificial seawater (ASW), (B) Mg<sup>2+</sup>-supplemented ASW (200 mM Mg<sup>2+</sup>), (C) 20 μM Gd<sup>3+</sup>-ASW, and (D) regular ASW.

mented ASW (200 mM Mg<sup>2+</sup>), 20 μM Gd<sup>3+</sup>-ASW, and regular ASW. Despite the discharge inhibition by Gd<sup>3+</sup> ions, R<sub>340/380</sub> transients were induced by the addition of K<sup>+</sup>-ASW (200 mM). The rise of R<sub>340/380</sub> measured in the presence of Gd<sup>3+</sup> ions was similar to that observed in Ca<sup>2+</sup>-free ASW and considerably lower than that observed in regular ASW [ $\Delta R_{340/380} = 0.23 \pm 0.11$ ,  $0.18 \pm 0.17$ , and  $0.62 \pm 0.24$ , respectively]. No significant difference was observed between Ca<sup>2+</sup>-free ASW ( $\Delta[Ca^{2+}]_i = 0.31 \pm 0.09$ ,  $n = 6$ ) and Mg<sup>2+</sup>-supplemented ASW ( $0.28 \pm 0.11$ ,  $n = 6$ ).

**Discussion**

*Ca<sup>2+</sup>-dependence of settlement and atrichous isorhiza discharge*

Although external Ca<sup>2+</sup> ions have been reported as being required for nematocyst discharge in several species of cnidaria (Salleo *et al.*, 1993; Santoro and Salleo, 1991a, b),

this is the first demonstration of Ca<sup>2+</sup>-dependent nematocyst discharge in larvae. We also confirmed and quantified the Ca<sup>2+</sup>-dependence of actinular settlement in *Tubularia mesembryanthemum*. Reduced-Ca<sup>2+</sup> seawater also inhibits metamorphosis of *Hydractinia echinata* planula larvae (Berking, 1988; Müller, 1985). The EC<sub>50</sub> of external Ca<sup>2+</sup> was 3 mM for *T. mesembryanthemum* actinular settlement, comparable to that of both *H. echinata* settlement and actinular AI discharge (Kawaii *et al.*, 1997). Since actinulae immersed in Ca<sup>2+</sup>-free ASW tended to disintegrate, settlement assays were not performed in Ca<sup>2+</sup>-free ASW.

The action of K<sup>+</sup>-ASW was strongly inhibited by increasing the Mg<sup>2+</sup> concentration of the bathing solution, and the inhibitory effects of the Mg<sup>2+</sup> ion increased when the external Ca<sup>2+</sup> concentrations were lowered. Similarly, Müller (1985) found that TPA-induced metamorphosis of *H. echinata* was promoted in Mg<sup>2+</sup>-reduced seawater. Mg<sup>2+</sup> and Ca<sup>2+</sup> ions compete with each other in many biological processes; consequently, we expected that an increase of Mg<sup>2+</sup> concentration in the bathing solution would reduce the Ca<sup>2+</sup> influx, thus inhibiting actinular settlement and AI discharge. However, the rise of R<sub>340/380</sub> values measured in Mg<sup>2+</sup>-supplemented ASW was equivalent to that observed in Ca<sup>2+</sup>-free ASW, and there was no influx of Ca<sup>2+</sup> ions from the Mg<sup>2+</sup>-supplemented ASW to nematocytes. These observations indicate that inhibition of AI discharge by Mg<sup>2+</sup> did not result from the lowering of nematocyte [Ca<sup>2+</sup>]<sub>i</sub> level by competitive influx of Mg<sup>2+</sup> and Ca<sup>2+</sup> ions. The enhancement of AI discharge and larval settlement by Mg<sup>2+</sup> reduction may be a result of cation-mediated alteration of membrane-associated cell function in signal transduction, and Mg<sup>2+</sup> ions would therefore regulate AI discharge and settlement of the hydroid as an inhibitory element.

*Involvement of stretch-activated (SA) channels*

It is natural for us to speculate about involvement of stretch-activated (SA) channels in actinular settlement, because the inhibitory effect of gadolinium ions at low concentrations in our study is comparable to that of opening SA channels in patch-clamped *Xenopus* oocytes (Yang and Sachs, 1989). Gd<sup>3+</sup> is the most effective blocker of SA channels found to date, although it also has some effect on Ca<sup>2+</sup> channels (Sadoshima *et al.*, 1992).

We previously demonstrated that both ATT contact with substrata and chemical stimuli are necessary to induce AI discharge (Kawaii *et al.*, 1997). By itself, a mechanical stimulus, such as immobilization of ATT by suction and vibration applied through a clean micropipette, did not trigger AI discharge. However, contact with a bacterial-film-coated micropipette did trigger discharge and also induced sinuous movement of the tentacles. Moreover, K<sup>+</sup>-ASW treatment did not result in AI discharge from ATTs

that were not in contact with any substrata. These results suggest that  $K^+$ -ASW treatment replaced sensory input of chemical stimuli, but did not mimic physical stimuli from ATT contact. Therefore, the observation that the stretching caused by immobilization of ATT did not trigger AI discharge does not rule out the involvement of SA channels in the discharge mechanism.

#### Effect of $Ca^{2+}$ -channel blockers

All the  $Ca^{2+}$ -channel blockers tested for inhibitory effects on  $K^+$ -ASW triggered AI discharge and larval settlement demonstrated similar dose-response curves and similar  $IC_{50}$  values. This consistency suggests that similar  $Ca^{2+}$ -channel types were involved in these events. Previously, we showed close relationships between discharge of AIs and sinuous movement of aboral tentacles (Kawaii *et al.*, 1997). Aboral tentacles that had discharged AIs usually initiated sinuous movements, resulting in settlement behavior. Thus AI discharge can be considered as the first step in the settlement process, and inhibition of AI discharge by  $Ca^{2+}$ -channel blockers interrupts the subsequent processes. Squashing of ATTs caused AI discharge, but no sinuous movement was observed when ASW was treated with a  $Ca^{2+}$ -channel blocker. These results suggest that these cations inhibited not only the AI discharge but also the signal transmission system that follows discharge and leads to the induction of sinuous movement or the aboral tentacle sinuous movement itself.

Direct monitoring of intracellular  $Ca^{2+}$  ions also indicated that AI discharge required both  $Ca^{2+}$ -release from intracellular stores and  $Ca^{2+}$ -influx from bathing solution, and that  $Ca^{2+}$ -channel blockers inhibited the large  $Ca^{2+}$  influx. Taking into consideration that  $K^+$ -ASW treatment triggered limited AI discharge in  $Ca^{2+}$ -free ASW (Kawaii *et al.*, 1997) or in ASW containing  $Ca^{2+}$ -channel blocker, a  $Ca^{2+}$  release from intracellular stores functioned as the signal transmission system in AI discharge. In the sea anemone *Haliplanella luciae*, a precise pharmacological study demonstrated that  $Ca^{2+}$  acts as a second messenger, and intracellular  $Ca^{2+}$  stores play an important role in the regulation of nematocyst discharge (Russell and Watson, 1995), via a mechanism of "calcium-induced calcium release" (Endo, 1977). It is conceivable that similar mechanisms for regulating nematocyst discharge are present in different types of nematocytes. In  $Ca^{2+}$ -free ASW or ASW containing  $10 \mu M Gd^{3+}$ ,  $K^+$ -ASW treatment triggered limited AI discharge and caused slight elevation of  $R_{340/380}$ . These small  $[Ca^{2+}]_i$  transients could be interpreted as a  $Ca^{2+}$  release from intracellular stores to nematocyte cytosol.

In conclusion, AI discharge was the primary action in actinular settlement, and signals from this discharge were spread throughout the larva, initiating actinular metamor-

phosis. The inhibitory effect of the  $Ca^{2+}$ -channel blockers demonstrated that AI discharge and subsequent signal transmission require involvement of  $Ca^{2+}$  channels.

#### Acknowledgments

The authors thank Dr. E. Hunter (The Centre for Environment, Fisheries & Aquaculture Science, Suffolk, U. K.) and Dr. C. G. Satuito (JAPAN NUS Co., Ltd., Tokyo, Japan) for their valuable suggestions.

#### Literature Cited

- Berking, S. 1988. Ammonia, tetraethylammonium, barium, and amiloride induce metamorphosis in the marine hydroid *Hydractinia*. *Roux's Arch. Dev. Biol.* **197**: 1–9.
- Endo, M. 1977. Calcium release from the sarcoplasmic reticulum. *Physiol. Rev.* **57**: 71–108.
- Freeman, G., and E. B. Ridgway. 1987. Endogenous photoprotein, calcium channels and calcium transients during metamorphosis in hydrozoans. *Roux's Arch. Dev. Biol.* **196**: 30–50.
- Freeman, G., and E. B. Ridgway. 1990. Cellular and intracellular pathways mediating the metamorphic stimulus in hydrozoan planulae. *Roux's Arch. Dev. Biol.* **199**: 63–79.
- Gitter, A. II., D. Oliver, and U. Thurm. 1994. Calcium- and voltage-dependence of nematocyst discharge in *Hydra vulgaris*. *J. Comp. Physiol. A* **175**: 115–122.
- Kawaii, S., K. Yamashita, M. Nakai, and N. Fusetani. 1997. Intracellular calcium transients during nematocyst discharge in actinulae of the hydroid, *Tubularia mesembryanthemum*. *J. Exp. Zool.* **278**: 299–307.
- McKay, M. C., and P. A. V. Anderson. 1988. Preparation and properties of cnidocytes from the sea anemone *Anthopleura elegantissima*. *Biol. Bull.* **174**: 47–53.
- Müller, W. A. 1985. Tumor-promoting phorbol esters induce metamorphosis and multiple head formation in the hydroid *Hydractinia*. *Differentiation* **29**: 216–222.
- Pantin, C. F. A. 1942. The excitation of nematocysts. *J. Exp. Biol.* **19**: 294–310.
- Russell, T. J., and G. M. Watson. 1995. Evidence for intracellular stores of calcium ions involved in regulating nematocyst discharge. *J. Exp. Zool.* **273**: 175–185.
- Sadoshima, J., T. Takahashi, L. Jahan, and S. Izumo. 1992. Roles of mechanosensitive ion channels, cytoskeleton, and contractile activity in stretch-induced immediate-early gene expression and hypertrophy of cardiac myocytes. *Proc. Natl. Acad. Sci. USA* **89**: 9905–9909.
- Salleo, A., G. Santoro, and P. Barra. 1993. Spread of experimentally induced discharge of the nematocytes in acontia of *Calliactis parasitica*. *Comp. Biochem. Physiol.* **104A**: 565–574.
- Salleo, A., G. La Spada, and R. Barbera. 1994a. Gadolinium is a powerful blocker of the activation of nematocytes of *Pelagia noctiluca*. *J. Exp. Biol.* **187**: 201–206.
- Salleo, A., G. La Spada, M. Drago, and G. Curcio. 1994b. Hyposmotic shock-induced discharge in acontia of *Calliactis parasitica* is blocked by gadolinium. *Experientia* **50**: 148–152.
- Santoro, G., and A. Salleo. 1991a. Cell-to-cell transmission in the activation of *in situ* nematocytes in acontia of *Calliactis parasitica*. *Experientia* **47**: 701–703.
- Santoro, G., and A. Salleo. 1991b. The discharge of *in situ* nematocysts of the acontia of *Aiptasia mutabilis* is a  $Ca^{2+}$ -induced response. *J. Exp. Biol.* **156**: 173–185.
- Tardent, P. 1988. History and current state of knowledge concerning the discharge of cnidae. Pp. 309–332 in *The Biology of Nematocysts*, D. A. Hessinger and H. M. Lenhoff eds. Academic Press, San Diego.

- Thorington, G. U., and D. A. Hessinger. 1988a.** Control of cnida discharge. I. Evidence for two classes of chemoreceptors. *Biol. Bull.* **174**: 163–171.
- Thorington, G. U., and D. A. Hessinger. 1988b.** Control of discharge: factors affecting discharge of cnidae. Pp. 233–253 in *The Biology of Nematocysts*, D. A. Hessinger and H. M. Lenhoff, eds. Academic Press, San Diego.
- Watson, G. M., and D. A. Hessinger. 1989.** Cnidocyte mechanoreceptors are tuned to the movements of swimming prey by chemoreceptors. *Science* **243**: 1589–1591.
- Watson, G. M., and D. A. Hessinger. 1991.** Chemoreceptor-mediated elongation of stereocilium bundles tunes vibration-sensitive mechanoreceptors on cnidocyte-supporting cell complexes to lower frequencies. *J. Cell Sci.* **99**: 307–316.
- Yamashita, K., S. Kawaii, M. Nakai, and N. Fusetani. 1997.** Behaviour and settlement of actinula larvae of *Tubularia mesembryanthemum* Allman, 1871 (Hydrozoa: Tubulariidae). Pp. 512–516 in *The Proceedings of the 6th ICCB*, J. C. den Hartog, ed. National Museum of Natural History, Leiden.
- Yanagita, T. M. 1973.** The 'cnidoblast' as an excitable system. *Publ. Seto Mar. Biol. Lab.* **20**: 675–693.
- Yang, X. -C., and F. Sachs. 1989.** Block of stretch-activated ion channels in *Xenopus* oocytes by gadolinium and calcium ions. *Science* **243**: 1068–1071.